

# COMPARISON BETWEEN PREDICTED AND EMPIRICAL $\Delta V_{\text{HB}}^{\text{Bump}}$ IN GALACTIC GLOBULAR CLUSTERS

M. ZOCCALI<sup>1</sup>, S. CASSISI<sup>2</sup>, G. PIOTTO<sup>1</sup>, G. BONO<sup>3</sup>, M. SALARIS<sup>4</sup>

*Draft version February 1, 2008*

## ABSTRACT

We present observational estimates of  $\Delta V_{\text{HB}}^{\text{Bump}}$  in a sample of 28 Galactic Globular Clusters (GGCs) observed by HST. The photometric accuracy and the sizable number of stars measured in each cluster allowed us to single out the RGB Bump both in metal-poor and in metal-rich GGCs.

Empirical values are compared with homogeneous theoretical predictions which account for both H and He burning phases over a wide range of metal abundances ( $0.0001 < Z < 0.02$ ). We found that, within current observational uncertainties on both iron and  $\alpha$ -element abundances, theory and observations are in very good agreement, provided that the metallicity scale by Carretta & Gratton (1997) as extended by Cohen et al. (1999) is adopted. Interestingly enough, we also found that both theoretical and observed values show a change in the slope of the  $\Delta V_{\text{HB}}^{\text{Bump}} - [\text{M}/\text{H}]$  relation toward higher metal contents.

*Subject headings:* Globular Clusters: general — stars: evolution — stars: horizontal branch — stars: Population II

## 1. INTRODUCTION

The RGB Bump is an intrinsic feature of the RGB Luminosity Function (LF) of GGCs. It appears as a peak in the differential LF or as a change in the slope of the cumulative LF. The presence of the Bump is due to the fact that during the RGB evolution the H-burning shell crosses the chemical discontinuity left over by the convective envelope soon after the first dredge-up phase. Since its first detection in 47 Tuc (King, Da Costa & Demarque 1985), it became the crossroad of several theoretical and observational investigations (Alves & Sarajedini 1999). The detection of the RGB Bump was mainly hampered by the size of the available samples of RGB stars. This is particularly true for the most metal-poor clusters, where the Bump moves toward brighter magnitudes and therefore less populated RGB regions. Only recently was this feature firmly detected in a large set of both Galactic (FP; Brocato et al. 1996) and extragalactic stellar systems (Desidera 1999).

The analysis by FP showed that the  $\Delta V_{\text{HB}}^{\text{Bump}}$  values predicted by theory were 0.4 mag brighter than the empirical estimates in a sample of 11 GGCs. This observable, which does not depend on the distance modulus, on the reddening, or on the calibration of photometric data, is a key parameter for assessing both the accuracy and the plausibility of the physical assumptions adopted for constructing the evolutionary models. As a consequence the discrepancy found by FP needs to be understood. In order to explain this mismatch between theory and observations Alongi et al. (1991) suggested the inclusion of convective undershooting at the boundary between the convective envelope and the thin H-burning shell, whereas Straniero,

Chieffi & Salaris (1992) called attention on the role that the global metallicity could play on the estimate of this observable. The latter hypothesis was confirmed by CS who found that standard models agree quite well with the observed  $\Delta V_{\text{HB}}^{\text{Bump}}$  values in 8 GGCs for which high resolution spectroscopic determinations of both  $[\text{Fe}/\text{H}]$  and  $[\alpha/\text{Fe}]$  were available. The significance of this result was limited by the small number of clusters taken into account and by the small metallicity range that they cover.

The main aims of this investigation are to provide new homogeneous measurements of both the Bump position and the corresponding  $\Delta V_{\text{HB}}^{\text{Bump}}$  value in a large sample of GGCs covering a wide metallicity range ( $-2.1 \leq [\text{Fe}/\text{H}] \leq -0.2$ ); and also to compare the new measurements with the predictions of standard evolutionary models. At present, the most homogeneous photometric sample comes from HST. Two of us (GP and MZ) are involved in a program aimed at collecting F439W and F555W WFPC2 images for all the GGC with  $(m - M)_B < 18.0$  not yet observed with HST. Ten of these clusters have been observed during Cycle 6 (GO6095) and nine more during the ongoing Cycle 7 (GP7470). Similar observations for nine additional clusters are available on the HST archive.

In §2 we present this sample of 28 GGCs, and discuss the data reduction strategy as well as the approach adopted for estimating both the zero age horizontal branch (ZAHB) and the RGB Bump visual magnitudes. Section 3 deals with the comparison between theory and observations, while in §4 we briefly summarize the main results.

## 2. DATA SELECTION AND REDUCTION

<sup>1</sup>Dipartimento di Astronomia, Università di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy; manu@pd.astro.it piotto@pd.astro.it

<sup>2</sup>Osservatorio Astronomico di Collurania, Via M. Maggini, 64100 Teramo, cassisi@astr.te.astro.it

<sup>3</sup>Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monteporzio Catone, Italy; bono@coma.mporzio.astro.it

<sup>4</sup>Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead L41 1LD, UK; ms@staru1.livjm.ac.uk

<sup>5</sup>The parameter  $\Delta V_{\text{HB}}^{\text{Bump}}$  is commonly defined as the difference in magnitude between the Red Giant Branch (RGB) Bump and the Horizontal Branch (HB) stars located within the RR Lyrae instability strip (see for a detailed discussion Fusi Pecci et al. 1990, hereinafter FP, and Cassisi & Salaris 1997, hereinafter CS).

The center of all the clusters were observed in the HST B (F439W) and V (F555W) bands with the WFPC2. Both very short (a few seconds) and long (a few hundreds of seconds) exposures are available for both filters allowing us to map the evolved regions of the color-magnitude diagram (CMD). The pre-processing, photometric reduction, and calibration to a standard B and V system of the data for each cluster were carried out following the same procedure as described in details in Piotto et al. (1999a) for a subset of images of the GO7470. The photometry for all the clusters extends from the tip of the RGB down to about 2 magnitudes below the turnoff, and include a number of stars ranging from  $\sim 5000$  for the loosest cluster to  $\sim 30000$  for the most concentrated ones. A detailed description of the CMDs will be presented in Piotto et al. (1999b). The large sample of stars, coupled with the high photometric accuracy, allows us to identify of the RGB Bump even in the most metal-poor clusters.

Figure 1 shows an example of the Bump identification in NGC5824, a metal-poor cluster in our database. The upper left panel shows the RGB differential LF. We constructed a histogram with a fixed bin size of 0.15 magnitude (*dotted line*) as well as the multibin histogram (*solid line*) described in Piotto et al. (1999c). The Bump is marked by a vertical arrow. As a further check of the accuracy in the Bump identification, the lower left panel shows the RGB cumulative LF. The Bump can be easily located since it is marked by a change in the slope. The right panel shows the CMD of NGC5824 and the arrow marks the position of the Bump along the RGB.

The determination of the ZAHB magnitude ( $V_{\text{ZAHB}}$ ) is a thorny problem, in particular for those clusters which show blue HBs. A standard method to determine the ZAHB magnitude for both intermediate and metal-poor clusters is to adopt the mean magnitude of the cluster RR Lyrae stars. Unfortunately, we could not adopt this method directly, since our photometry covers a very short time interval, and therefore the RR Lyrae were always measured at random pulsation phases (Piotto et al. 1999a).

In order to overcome this problem we have undertaken a different approach. For metal-poor and intermediate metallicity clusters we selected from the literature three clusters which offer accurate photometry, a large number of RR Lyrae stars, and a well defined blue HB tail. The template clusters are the following: NGC1851 (Walker 1998) for clusters in the metallicity range  $-1.5 < [\text{Fe}/\text{H}] < -1.0$ , NGC5272 (M3, Buonanno et al. 1994) for clusters in the range  $-1.7 < [\text{Fe}/\text{H}] < -1.5$ , and NGC4590 (M68, Walker 1994) for more metal-poor clusters. The CMDs of the template clusters were artificially shifted in color and in magnitude in order to match both the RGB and the blue HB tail of each cluster. The mean RR Lyrae magnitude of the template cluster was scaled according to the magnitude shift adopted to overlap the stellar distributions along the HB. The  $V_{\text{ZAHB}}$  magnitudes were estimated by using the relation between the mean RR Lyrae magnitude and the  $V_{\text{ZAHB}}$  magnitude suggested by CS. It is worth noting that this method is totally independent from any zero point difference between the template photometry and ours.

For the most metal-rich clusters ( $[\text{Fe}/\text{H}] > -1$ ) with well

populated red HBs but no RR Lyraes, first we estimated the photometric error  $\sigma_V^2 = \sigma_{(B-V)}^2/2$ , where  $\sigma_{(B-V)}^2$  is the standard deviation of the color distribution of the RGB stars at the level of the HB. Then we fixed  $V_{\text{ZAHB}}$  at  $3\sigma_V$  magnitudes above the lower envelope of HB stellar distribution. However, in order to provide a consistent determination of this parameter over the whole GGCs sample, the  $V_{\text{ZAHB}}$  at the level of RR Lyrae instability strip was evaluated according to the method suggested by Fullton et al. (1995). Figure 1 shows the fit of the same metal-poor cluster NGC5824, (full dots) with the template cluster M68 (crosses). A vertical and a horizontal shift of 2.83 and 0.08 magnitudes were applied to M68 for matching both the RGB and the HB. The M68 RR Lyrae stars are plotted as open squares. Taking into account the uncertainties in the fit, we estimated an error in  $V_{\text{ZAHB}}$  of the order of 0.1 mag.

Table 1 summarizes the cluster observables: column (1) gives the NGC and the Messier numbers; column (2) lists the visual magnitudes of the RGB Bump and the photometric error; column (3) gives the mean RR Lyrae visual magnitudes estimated according to the method previously described; column (4) lists the ZAHB visual magnitude at the lower envelope of red HB stars in metal-rich clusters; column (5) gives the cluster metallicities according to the Carretta & Gratton (1997, hereinafter CG) scale<sup>6</sup>.

### 3. COMPARISON BETWEEN THEORY AND OBSERVATIONS

Figure 2 shows the comparison between theory and observations in the  $[\text{M}/\text{H}]-\Delta V_{\text{HB}}^{\text{Bump}}$  plane. The global metallicities were estimated by adopting a mean  $\alpha$  enhancement equal to 0.3 for clusters with  $[\text{Fe}/\text{H}] < -1.0$  and to 0.20 for more metal-rich clusters. The former value was suggested by Carney (1996, hereinafter C96), while the latter, due to the paucity of data available in the literature, is a mean between the estimates collected by C96 and by Salaris & Cassisi (1996, hereinafter SC). The observed  $\Delta V_{\text{HB}}^{\text{Bump}}$  values have been plotted in the top panel according to the CG metallicity scale. The  $\Delta V_{\text{HB}}^{\text{Bump}}$  error bars have been calculated by quadratically combining the errors on  $V_{\text{ZAHB}}$  and on  $V_{\text{Bump}}$ . The global metallicity error bars are a lower limit of the uncertainties affecting both  $[\text{Fe}/\text{H}]$  and  $[\alpha/\text{Fe}]$  measurements (see C96 and Rutledge, Hesser, & Stetson 1998, hereinafter RHS).

The theoretical predictions plotted in the top panel were estimated by adopting progenitor masses ranging from  $M/M_{\odot} = 0.8$  to 1.0 and a wide range of global metallicities ( $-2.3 \leq [\text{M}/\text{H}] \leq 0.0$ ). The initial helium contents adopted in constructing evolutionary models are the following:  $Y = 0.23$  for  $[\text{M}/\text{H}] \leq -0.5$ ,  $Y = 0.255$  for  $[\text{M}/\text{H}] = -0.25$ , and  $Y = 0.289$  for  $[\text{M}/\text{H}] = 0.0$ . Metal-poor  $\Delta V_{\text{HB}}^{\text{Bump}}$  theoretical estimates up to  $[\text{M}/\text{H}] \approx -0.5$  already been presented in CS, whereas more metal-rich ones were specifically computed, to extend the predictions to clusters more metal-rich than 47 Tuc. Basic assumptions on the input physics adopted for constructing evolutionary models were extensively described in CS and Bono et al. (1997 and references therein) and therefore they are not

<sup>6</sup>For those clusters whose metallicity was not provided by CG, the cluster metallicities collected by Harris (1996) were transformed into the CG scale by adopting the new calibration provided by Cohen et al. (1999, hereinafter CGBC). In contrast with CG the new scale applies also to metal-rich clusters ( $-2.12 \leq [\text{Fe}/\text{H}] \leq -0.3$ ).

discussed here. The reader interested in a detailed discussion on the dependence of  $\Delta V_{\text{HB}}^{\text{Bump}}$  on mixing-length, He content, and element diffusion is referred to CS and to Cassisi, Degl'Innocenti, & Salaris (1997, hereinafter CDS). Bolometric magnitudes were transformed into V magnitudes by adopting bolometric corrections provided by Castelli, Gratton & Kurucz (1997).

In order to account for the  $V_{\text{Bump}}$  dependence on cluster age as suggested by CS and more recently by Alves & Sarajedini (1999) in a detailed investigation on HST data of 8 SMC clusters, we plotted the theoretical predictions for three different ages: 12 (short-dash line), 14 (solid line), and 16 (long-dash line) Gyr. In the last few years a large number of theoretical and observational investigations have been devoted to the absolute and the relative ages of GGCs as well as to the errors affecting such parameters. In fact, they depend on the physical assumptions and on the input physics (Cassisi et al. 1998; Vandenberg, Stetson & Bolte 1996) adopted for constructing evolutionary models. As plausible assumptions we adopted an average cluster age of 14 Gyr (Vandenberg 1999) and an average uncertainty of  $\pm 2$  Gyr. The change in the slope toward higher metal contents shown by theoretical predictions is due to the fact that at fixed age an increase in the evolving mass causes a smoother increase in the core-mass luminosity relation, and in turn in the  $V_{\text{Bump}}$  magnitude. In fact, for metallicities ranging from  $Z=0.001$  to  $Z=0.006$  the  $M_V(\text{Bump})$  and the  $M_V(\text{ZAHB})$  magnitudes changes according to the following derivatives:  $\partial M_V(\text{ZAHB})/\partial [\text{M}/\text{H}] \approx 0.093$  and  $\partial M_V(\text{Bump})/\partial [\text{M}/\text{H}] \approx 0.753$ , whereas for  $0.006 \leq Z \leq 0.02$  they change according to  $\approx 0.289$  and  $\approx 1.251$  respectively.

The data plotted in the top panel of Figure 2 show clearly that the discrepancy of 0.4 magnitudes suggested by FP is completely removed over the entire metallicity range, and indeed the trend of empirical data is well reproduced by standard models. This result is even more compelling if we take into account that the previous comparisons (FP and CS) were hampered by the small number of metal-poor clusters ( $[\text{Fe}/\text{H}] < -1.5$ ) for which reliable estimates of  $V_{\text{Bump}}$  were available. Figure 2 also extends the comparison to  $[\text{Fe}/\text{H}] = -0.2$ ; note how the observed values in the high metallicity range show the flattening predicted by the theory, though with a distribution that is somehow flatter. A plausible change in the He content (CS) and/or the inclusion of element diffusion (CDS) can account for this effect only marginally. As a consequence, this result suggests that the metal-rich clusters could be younger than the bulk of our clusters (Salaris & Weiss 1998; Rosenberg et al. 1999).

In the cluster sample adopted in this investigation there are only two clusters -NGC7078 and NGC5694- which are marginally in agreement with the theoretical expectations. For both clusters the uncertainty on the location of the Bump is very small (Table 1). In the case of NGC7078, the adopted mean  $V_{\text{RR}}$  magnitude is also in very good agreement with the value obtained by Silberman & Smith (1995). Therefore for this cluster the discrepancy could be due to an underestimate of the cluster metallicity and/or of the  $\alpha$  enhancement (see e.g. CG). An independent estimate of  $V_{\text{RR}}$  for NGC5694 is not available, but any plausi-

ble assumption on its uncertainty can hardly remove such a discrepancy.

In order to account for the uncertainty on the metallicity scale the middle panel shows the same comparison as the top panel, but the observed points are plotted according to the Zinn & West (1984, hereinafter ZW) metallicity scale. The data plotted in this panel show that the  $\Delta V_{\text{HB}}^{\text{Bump}}$  values are systematically shifted toward lower metal contents when compared with theoretical observables. At present, both systematic and observational errors affecting the metallicity ranking of GGCs are still controversial issues (RHS; C96). This notwithstanding, the CG scale is more robust since it relies on recent high dispersion spectroscopic measurements and up-to-date atmosphere models. Even though current observational uncertainties affects the global metallicity of individual clusters, data plotted in Figure 2 support the evidence that the ZW scale underestimates the cluster metallicity in the range  $-1.7 < [\text{Fe}/\text{H}] < -1.0$ . In order to supply a quantitative estimate of the difference we performed a fit of the empirical data ( $-2.0 < [\text{Fe}/\text{H}] < -0.5$ ) with predictions at 14 Gyr. The standard deviation is 0.05 mag for the CGBC scale and 0.17 mag for the ZW scale. The latter value is almost a factor of two larger than the photometric uncertainty.

#### 4. CONCLUSIONS

We have presented new homogeneous measurements of the  $\Delta V_{\text{HB}}^{\text{Bump}}$  values for a sample of 28 GGCs observed with HST, and a detailed comparison with the theoretical models. By relying on homogeneous theoretical and observational frameworks and on the metallicity scale suggested by CG and by CGBC we found that, within current uncertainties, observables predicted by standard H- and He-burning evolutionary models agree with the empirical data. This result is further strengthened by the fact that this comparison was extended from metal-poor to metal-rich clusters ( $-2.1 \leq [\text{Fe}/\text{H}] \leq -0.2$ ).

New theoretical predictions for metal-rich clusters show a change of the slope of the  $\Delta V_{\text{HB}}^{\text{Bump}} - [\text{M}/\text{H}]$  relation. This behavior is supported by the tail of metal-rich clusters in our sample and does not depend on the adopted metallicity scale. Leading physical arguments on the dependence of  $\Delta V_{\text{HB}}^{\text{Bump}}$  on input physics support the suggestion that metal-rich cluster could be younger than the bulk of clusters in our sample.

By adopting the ZW metallicity scale we found that empirical data at low and intermediate metallicity are shifted toward lower metallicities when compared with theory. At the same time the agreement between theory and observations supports the use of a  $\Delta V_{\text{HB}}^{\text{Bump}}$  - metallicity relation for constraining the cluster metallicity (Desidera 1999). However, we note that such a relation relies on the assumption that all GGCs are coeval within  $\pm 1$  Gyr (Stetson et al. 1999), and that the intrinsic accuracy is of the order of 0.15 dex provided that the  $\Delta V_{\text{HB}}^{\text{Bump}}$  values are measured with an accuracy of 0.10 mag.

We are deeply indebted to E. Carretta and R. Gratton for providing us with the extension of the CG metallicity scale to  $[\text{Fe}/\text{H}] = -0.3$  in advance of publication.

## REFERENCES

- Alongi, M., Bertelli, G., Bressan, A. & Chiosi, C. 1991, A&A, 244, 95
- Alves, D. R., & Sarajedini, A. 1999, ApJ, 511, 225
- Bono, G., Caputo, F., Cassisi, S., Castellani, V., & Marconi, M. 1997, ApJ, 479, 279.
- Brocato, E., Buonanno, R., Malakhova, Y., & Piersimoni, A. M. 1996, A&A, 311, 778
- Buonanno, R., Corsi, C. E., Buzzoni, A., Cacciari, C., Ferraro, F. R., & Fusi Pecci, F. 1994, A&A, 290, 69
- Carney, B. W. 1996, PASP, 108, 900
- Carretta E., & Gratton, R. G. 1996, A&AS, 121, 95 (CG)
- Cassisi, S., Castellani, V., Degl'Innocenti, S., & Weiss, A. 1998, A&AS, 129, 267 (CDS)
- Cassisi, S., Degl'Innocenti, S., & Salaris, M. 1997, MNRAS, 290, 515
- Cassisi, S., & Salaris, M. 1997, MNRAS, 285, 593 (CS)
- Castelli, F., Gratton, R. G., & Kurucz, R. 1997, A&A, 318, 841
- Cohen, J. G., Gratton, R. G., Behr, B. B., & Carretta, E. 1999, in preparation (CGBC)
- Desidera, S. 1999, in ESO Workshop, Chemical Evolution from Zero to High Redshift, in press
- Fullton, L. K., Carney, B. W., Olzewski, E. W., Zinn, R., Demarque, P., Janes, K. A., Da Costa, G. S. & Seitzer, P. 1995, AJ, 110, 652
- Fusi Pecci, F., Ferraro, F. R., Crocker, D. A., Rood, R. T., & Buonanno, R. 1990, A&A, 238, 95 (FP)
- Harris, W. E. 1996, AJ, 112, 1487
- King, C. R., Da Costa, G.S., & Demarque, P. 1985, ApJ, 299, 674
- Piotto, G. et al. 1999a, AJ, 117, 264
- Piotto, G. et al. 1999b, in preparation
- Piotto, G. et al. 1999c, AJ, submitted
- Rosenberg, A., Saviane, I., Piotto, G., & Aparicio, A. 1999, in preparation
- Rutledge, G. A., Hesser, J. E., & Stetson, P. B. 1997, 109, 907 (RHS)
- Salaris, M., & Cassisi, S. 1996, A&A, 305, 858 (SC)
- Salaris, M., & Weiss, A. 1998, A&A, 335, 943
- Silberman, N. A., & Smith, H. 1995, AJ, 110, 704
- Stetson, P. B., et al. 1999, AJ, 117, 247
- Straniero, O., Chieffi, A., & Salaris, M. 1992, Mem. SAI., 63, 315
- Vandenberg, D. A. 1999, in The Third Stromlo Symposium: The Galactic Halo, ed. B.K. Gibson, T.S. Axelrod & M.E. Putnam (San Francisco: ASP), in press
- Vandenberg, D. A., Stetson, P. B., & Bolte, M. 1996, ARA&A, 34, 461
- Walker, A. R. 1994, AJ, 108, 555
- Walker, A. R. 1998, AJ, 116, 220
- Zinn, R., & West, M. J. 1984, ApJS, 55, 45 (ZW)

TABLE 1  
CLUSTER OBSERVABLES

| Object<br>(1) |       | $V_{\text{Bump}}$<br>(2) | $V_{\text{RR}}$<br>(3) | $V_{\text{RHB}}$<br>(4) | $[Fe/H]_{CG}$<br>(5) |
|---------------|-------|--------------------------|------------------------|-------------------------|----------------------|
| 104           | 47Tuc | $14.57 \pm 0.02$         | ...                    | 14.16                   | $-0.70^a$            |
| 362           |       | $15.47 \pm 0.02$         | 15.33                  | ...                     | $-1.15^a$            |
| 1851          |       | $16.16 \pm 0.02$         | 16.05                  | ...                     | $-1.14^b$            |
| 1904          | M79   | $16.00 \pm 0.04$         | 16.21                  | ...                     | $-1.37^a$            |
| 2808          |       | $16.31 \pm 0.03$         | 16.25                  | ...                     | $-1.24^b$            |
| 5634          |       | $17.77 \pm 0.03$         | 17.95                  | ...                     | $-1.66^b$            |
| 5694          |       | $18.49 \pm 0.04$         | 18.64                  | ...                     | $-1.70^b$            |
| 5824          |       | $18.10 \pm 0.03$         | 18.46                  | ...                     | $-1.69^b$            |
| 5927          |       | $17.37 \pm 0.04$         | ...                    | 16.83                   | $-0.31^b$            |
| 6093          | M80   | $16.12 \pm 0.03$         | 16.36                  | ...                     | $-1.48^b$            |
| 6139          |       | $18.30 \pm 0.03$         | 18.40                  | ...                     | $-1.50^b$            |
| 6205          | M13   | $14.70 \pm 0.04$         | 14.99                  | ...                     | $-1.39^a$            |
| 6235          |       | $17.24 \pm 0.05$         | 17.31                  | ...                     | $-1.27^b$            |
| 6273          | M19   | $16.77 \pm 0.04$         | 16.91                  | ...                     | $-1.53^b$            |
| 6284          |       | $17.36 \pm 0.05$         | 17.25                  | ...                     | $-1.20^b$            |
| 6287          |       | $16.60 \pm 0.05$         | 17.21                  | ...                     | $-1.88^b$            |
| 6293          |       | $16.04 \pm 0.05$         | 16.47                  | ...                     | $-1.76^b$            |
| 6342          |       | $18.00 \pm 0.05$         | ...                    | 17.56                   | $-0.57^b$            |
| 6356          |       | $18.53 \pm 0.03$         | ...                    | 18.05                   | $-0.44^b$            |
| 6362          |       | $15.60 \pm 0.02$         | 15.35                  | ...                     | $-0.96^a$            |
| 6388          |       | $17.69 \pm 0.04$         | ...                    | 17.31                   | $-0.53^b$            |
| 6441          |       | $18.46 \pm 0.04$         | ...                    | 17.99                   | $-0.46^b$            |
| 6522          |       | $17.01 \pm 0.04$         | 17.06                  | ...                     | $-1.38^b$            |
| 6624          |       | $16.68 \pm 0.02$         | ...                    | 16.19                   | $-0.36^b$            |
| 6652          |       | $16.44 \pm 0.02$         | ...                    | 16.11                   | $-0.86^b$            |
| 6934          |       | $16.85 \pm 0.03$         | 17.01                  | ...                     | $-1.40^b$            |
| 6981          | M72   | $17.13 \pm 0.04$         | 17.26                  | ...                     | $-1.40^b$            |
| 7078          | M15   | $15.41 \pm 0.04$         | 15.86                  | ...                     | $-2.12^a$            |

<sup>a</sup>CG metallicities from high resolution spectroscopy.

<sup>b</sup>Cluster metallicities based on CGBC scale.




Fig. 1.— Example of the empirical measure of the Bump and ZAHB location. The arrows indicate the adopted Bump magnitude. *Panel a:* Differential LF; *Panel b:* cumulative LF; *Panel c:* position of the Bump in the CMD; *Panel d:* match between the HBs of NGC5824 and of the template cluster M68.




Fig. 2.— Comparison between the theoretical and empirical values of  $\Delta V_{\text{HB}}^{\text{Bump}}$  as a function of the global metallicity. *Top panel:* the empirical data plotted according to the CG metallicity scale (see text for more details). *Bottom panel:* Same as the top panel but with the empirical data were plotted according to the ZW metallicity scale.